

Determination of the Median Effective Concentration of Nano-Zero Valent Iron to the Marine Phytoplankton Species *Isochrysis galbana*

Ahmad Razi Othman^{1*}

¹Research Centre for Sustainable Process Technology (CESPRO), Faculty of Engineering and Built Environment, Universiti Kebangsaan Malaysia, 43600 UKM Bangi, Selangor, Malaysia.

*Corresponding author:

Ahmad Razi Othman

Research Centre for Sustainable Process Technology (CESPRO),
Faculty of Engineering and Built Environment,
Universiti Kebangsaan Malaysia,
43600 UKM Bangi,
Selangor,
Malaysia.

Email: ahmadrazi@ukm.edu.my

HISTORY

Received: 11th Oct 2018
Received in revised form: 27th of Nov 2018
Accepted: 25th of Dec 2018

KEYWORDS

four-parameter logistics
toxicity
EC₅₀
Isochrysis galbana
nano zero valent iron

ABSTRACT

Nano zero valent iron (nZVI) is often used in the remediation of organic and inorganic contaminant and has been hailed as an emerging remediation tool. Unfortunately, few studies have been done on the toxicity of nZVI on organisms especially aquatic organisms. In this study, the toxicity of commercial forms (uncoated, organic coating, and iron oxide coating) of nano zero-valent iron (nZVI) to the marine phytoplankton *Isochrysis galbana* is revisited in terms of the determination of the EC₅₀ or median effective concentration using nonlinear regression. Based on the four-parameter logistics equation, the median effective concentration for Nanofer 25S, Fe²⁺ and Fe³⁺ were 5.46 mg/L (95% confidence interval of 4.84 to 6.16), 70.89 mg/L (95% confidence interval of 37.26 to 134.9), 64.29 mg/L (95% confidence interval of 44.92 to 92.02). The results indicate that Nanofer 25S was the most toxic.

INTRODUCTION

Zero valent iron (ZVI or Fe⁰) is a reducing agent that commonly used in remediating contaminant from environment. This reducing agent is readily available, inexpensive and can remediate various contaminants [1]. Furthermore, ZVI is proven to be effective in reducing stable contaminants such as chlorinated hydrocarbon from the environment [2]. This strong reducing agent is proven to be effective in reducing various types of metals from waste water such as chromium (Cr (VI)), copper (Cu), molybdenum (Mo) and lead (Pb) [3–5]. The utilization of ZVI is not only focusing on the remediating metals, it is also reported to be effectively in remediating leachate from agricultural activities. Nitrate contamination originating from pesticide and herbicide used in agriculture is reported to contaminate clean water source [6].

A combination technology of bacteria associated with ZVI is proven to be better in nitrate removal from agriculture waste water when compared to bacteria treatment and ZVI treatment alone [7,8]. ZVI also can be used in the treatment of pharmaceutical waste. Almost 90% of amoxicillin content can be removed by the ZVI from the respected waste water [9]. Whilst ZVI is commonly used in pollutant treatment, it is also can be used in industrial process. ZVI is reported to enhanced the delignification process of palm oil empty fruit bunch into a fine

material that can be use later in biopolymer and biofuel industry [10]. This material also accelerate the synthesis of short chain fatty acid in the industry [11].

Despite of its beneficial property, ZVI treatment is also believed to contribute to the biota toxicity. The ZVI toxicity showed a vast effect towards wide range organism in the environment. This include effect towards microorganism, aquatic organism and plant [12]. Toxicity analysis towards phytoplankton population exposed to the ZVI was evaluated and the result shows a significant decreasing in the phytoplankton population [13]. Furthermore, utilization of ZVI in remediating pollution can effect microorganism population such as bacteria and archaea [14]. The contribution of ZVI toxicity is believed due to the formation of reactive oxygen species (ROS) formed through the Fenton reaction and Haber-Weiss reaction [15,16]. Other factor that contributes to ZVI toxicity is due to the accumulation of iron in the organism.

Accumulation of iron from ZVI contribute to malfunction in algae, plant and fish [17–22]. Lastly, accumulation of ZVI in environment reducing the dissolve oxygen (DO) amount in water body [20]. Due to the presence of toxicity characteristic in nZVI, it is important to determine the EC₅₀ or median effective concentration by nZVI towards microorganisms. The median effective concentration can be calculated using mathematical

software. An LD₅₀ plot is in non-linear plot where discrimination between the value can only be calculated precisely through statistic [23]. This statistical approach is used to minimize the lack-of-fit error caused by the improper mathematical modelling selection [24].

MATERIALS AND METHODS

Acquisition of data

Data from the works of Keller et al. [13] from figure five showing the growth rate of *I. galbana* exposed to the various types of ZVI. The information was processed by the Webplotdigitizer 2.5 software that digitizes the scanned figure into comma separated values.

Four-Parameter Logistics Modelling

The four-parameter logistics equation was chosen in plotting a non-linear regression based on the least square fitting [25]. The equation is described as follows;

$$y = Bottom + \frac{(Top - Bottom)}{1 + 10^{(Log EC_{50} - x) * Hillslope}}$$

where y represent the growth rate (cell/H) obtained, x is the concentration of nZVI exposed to the phytoplankton (log unit), $Log EC_{50}$ is the amount that produces 50% of signal response and Hillslope is the slope-like parameter (Hill coefficient). Multivariate analysis using the four-parameter supply model was calculated using the PRISM software (5.0) accessible from www.graphpad.com. Growth rate for *I. galbana* exposed to Fe^{2+} showed no inhibition trend and was excluded from the remodeling.

RESULTS AND DISCUSSION

The results of the modelling exercise show acceptable fitting of the data to the model (Figs 1 to 3). Based on the four-parameter logistics equation, the median effective concentration for Nanofer 25S, Fe^{2+} and Fe^{3+} were 5.46 mg/L (95% confidence interval of 4.84 to 6.16), 70.89 mg/L (95% confidence interval of 37.26 to 134.9), 64.29 mg/L (95% confidence interval of 44.92 to 92.02). The coefficient of determination or R^2 were 0.99, 0.78 and 0.88, respectively, which indicate that more data points are needed for the two latter cases to improve determination reliability. Despite this setback, the results indicate that Nanofer 25S was the most toxic. In comparison to other published works, the median effective concentration for zerovalent iron nanoparticles against cyanobacteria is 50 mg/L whilst for *D. magna* is greater than 1000 mg/L [26].

The determination of median effective concentration for toxicant along with its confidence interval is an important exercise neglected by numerous researchers. The uncertainty values obtained can be used to compare the toxicity of various toxicants and target organisms to each other. The least information that can be obtained is comparative assessment across publications. Although the comparison of various toxicants to the same organisms or the comparison of various organisms to the same toxicant is best done within the same lab and time, in reality this is impossible or uneconomical to be carried out.

If comparison is to be made across publications, the confidence interval should be used as an assessment of statistical significance, whether one result is more toxic or sensitive to another. Significant difference between two values occurs when the 95% confidence interval values are non-overlapped.

However, when the 95% confidence interval values overlapped, this does not means significant or not significant at the $p < 0.05$ level. What overlapped confidence interval values means is that more data and study are needed [27,28].




Fig. 1. The effect of Nanofer 25S on the growth rate for *I. galbana*.




Fig. 2. The effect of Nanofer STAR on the growth rate for *I. galbana*.




Fig. 3. The effect of Fe^{3+} on the growth rate for *I. galbana*.

CONCLUSION

Nano zero valent iron (nZVI) is an emerging remediation tool. Its toxicity to aquatic organisms, however needs to be determined through the use of nonlinear regression technique. In this study, the toxicity of commercial forms (uncoated, organic coating, and iron oxide coating) of nano zero-valent iron (nZVI) to the marine phytoplankton *Isochrysis galbana* is revisited in terms of the

determination of the EC₅₀ or median effective concentration using the four-parameter logistics model. Based on the four-parameter logistics model, the results indicate that Nanofer 25S was the most toxic.

REFERENCES

- Li J, Dou X, Qin H, Sun Y, Yin D, Guan X. Characterization methods of zerovalent iron for water treatment and remediation. *Water Res.* 2019 Jan;148:70–85.
- Pramudita D, Iskandar I, Indarto A. Nano-Enhanced Materials for Reclamation of Mine Spoils. In: *Bio-Geotechnologies for Mine Site Rehabilitation*. Elsevier; 2018. p. 201–14.
- Qian L, Shang X, Zhang B, Zhang W, Su A, Chen Y, et al. Enhanced removal of Cr(VI) by silicon rich biochar-supported nanoscale zero-valent iron. *Chemosphere.* 2019 Jan 1;215:739–45.
- Vollprecht D, Krois L-M, Sedlazeck KP, Müller P, Mischitz R, Olbrich T, et al. Removal of critical metals from waste water by zero-valent iron. *J Clean Prod.* 2019 Jan 20;208:1409–20.
- Liu X, Lai D, Wang Y. Performance of Pb(II) removal by an activated carbon supported nanoscale zero-valent iron composite at ultralow iron content. *J Hazard Mater.* 2019 Jan 5;361:37–48.
- Okafor PN, Ogbonna UI. Nitrate and nitrite contamination of water sources and fruit juices marketed in South-Eastern Nigeria. *J Food Compos Anal.* 2003 Apr 1;16(2):213–8.
- Mofradnia SR, Ashouri R, Tavakoli Z, Shahmoradi F, Rashedi H, Yazdian F, et al. Effect of zero-valent iron/starch nanoparticle on nitrate removal using MD simulation. *Int J Biol Macromol.* 2019 Jan 1;121:727–33.
- Zhang Y, Douglas GB, Kaksonen AH, Cui L, Ye Z. Microbial reduction of nitrate in the presence of zero-valent iron. *Sci Total Environ.* 2019 Jan 1;646:1195–203.
- Pirsaheb M, Moradi S, Shahlaei M, Wang X, Farhadian N. A new composite of nano zero-valent iron encapsulated in carbon dots for oxidative removal of bio-refractory antibiotics from water. *J Clean Prod.* 2019 Feb 1;209:1523–32.
- Santanaraj J, Sajab MS, Mohammad AW, Harun S, Chia CH, Zakaria S, et al. Enhanced Delignification of Oil Palm Empty Fruit Bunch Fibers with in situ Fenton-oxidation. *BioResources.* 2017 Jun 6;12(3):5223–35.
- Luo J, Feng L, Chen Y, Li X, Chen H, Xiao N, et al. Stimulating short-chain fatty acids production from waste activated sludge by nano zero-valent iron. *J Biotechnol.* 2014 Oct 10;187:98–105.
- Lei C, Sun Y, Tsang DCW, Lin D. Environmental transformations and ecological effects of iron-based nanoparticles. *Environ Pollut.* 2018 Jan 1;232:10–30.
- Keller AA, Garner K, Miller RJ, Lenihan HS. Toxicity of Nano-Zero Valent Iron to Freshwater and Marine Organisms. Chin W-C, editor. *PLoS ONE.* 2012 Aug 30;7(8):e43983.
- Zabetakis KM, Niño de Guzmán GT, Torrents A, Yarwood S. Toxicity of zero-valent iron nanoparticles to a trichloroethylene-degrading groundwater microbial community. *J Environ Sci Health Part A.* 2015 Jul 3;50(8):794–805.
- Wang B, Yin J-J, Zhou X, Kurash I, Chai Z, Zhao Y, et al. Physicochemical Origin for Free Radical Generation of Iron Oxide Nanoparticles in Biomicroenvironment: Catalytic Activities Mediated by Surface Chemical States. *J Phys Chem C.* 2013 Jan 10;117(1):383–92.
- Wu H, Yin J-J, Wamer WG, Zeng M, Lo YM. Reactive oxygen species-related activities of nano-iron metal and nano-iron oxides. *J Food Drug Anal.* 2014 Mar 1;22(1):86–94.
- Toh PY, Ng BW, Chong CH, Ahmad AL, Yang J-W, Chieh Derek CJ, et al. Magnetophoretic separation of microalgae: the role of nanoparticles and polymer binder in harvesting biofuel. *RSC Adv.* 2014 Dec 16;4(8):4114–21.
- Japar AS, Takriff MS, Jahim JM, Kadhun AAH. Acetone–Butanol–Ethanol Fermentation From Palm Oil Mill Effluent Using *Clostridium acetobutylicum*. In: *Developments in Sustainable Chemical and Bioprocess Technology*. Boston, MA: Springer US; 2013. p. 35–41.
- Ma X, Gurung A, Deng Y. Phytotoxicity and uptake of nanoscale zero-valent iron (nZVI) by two plant species. *Sci Total Environ.* 2013 Jan 15;443:844–9.
- Chen P-J, Tan S-W, Wu W-L. Stabilization or Oxidation of Nanoscale Zerovalent Iron at Environmentally Relevant Exposure Changes Bioavailability and Toxicity in Medaka Fish. *Environ Sci Technol.* 2012 Aug 7;46(15):8431–9.
- Munro HN (Hamish N. Mammalian protein metabolism. Volume III. 591 p.
- Schiwy A, Maes HM, Koske D, Flecken M, Schmidt KR, Schell H, et al. The ecotoxic potential of a new zero-valent iron nanomaterial, designed for the elimination of halogenated pollutants, and its effect on reductive dechlorinating microbial communities. *Environ Pollut.* 2016 Sep 1;216:419–27.
- Shukor MY. Remodelling of the Calibration Curve for the Potentiometric Sensing of Lamotrigine Based on Molecularly Imprinted Polymers Using the Four-Parameter Logistic Model. *J Environ Microbiol Toxicol.* 2017;5(1):1–3.
- Alfaro ME, Zoller S, Lutzoni F. Bayes or Bootstrap? A Simulation Study Comparing the Performance of Bayesian Markov Chain Monte Carlo Sampling and Bootstrapping in Assessing Phylogenetic Confidence. *Mol Biol Evol.* 2003 Feb 1;20(2):255–66.
- Motulsky HJ, Ransnas LA. Fitting curves to data using nonlinear regression: a practical and nonmathematical review. *FASEB J Off Publ Fed Am Soc Exp Biol.* 1987 Nov 1;1(5):365–74.
- Marsalek B, Jancula D, Marsalkova E, Mashlan M, Safarova K, Tucek J, et al. Multimodal action and selective toxicity of zerovalent iron nanoparticles against cyanobacteria. *Environ Sci Technol.* 2012 Jan 31;46(4):2316–23.
- Schenker N, Gentleman JF. On judging the significance of differences by examining the overlap between confidence intervals. *Am Stat.* 2001;55(3):182–6.
- Asri NAM, Begum G, Ahmad SA, Yasid NA, Shukor MY. A confidence interval comparison chart to assess the sensitivity of acetylcholinesterase from *Hemibagrus nemurus* as a source of enzyme for insecticides detection. *J Environ Microbiol Toxicol.* 2017 Jul 31;5(1):14–20.