

# Determination of Supercritical Carbon Dioxide Extraction Parameters for *Quercus infectoria* Galls and the Effects on Extraction Yield and Antioxidant Activity

Liza Md Salleh<sup>a,b,\*</sup>, Hasmida Mohd Nasir<sup>a</sup>, Harisun Yaakob<sup>c</sup>, Mohd Azizi Che Yunus<sup>a</sup>

<sup>a</sup>Centre of Lipid Engineering and Applied Research (CLEAR), Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia

<sup>b</sup>Department of Bioprocess Engineering, Faculty of Chemical Engineering, Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia

<sup>c</sup>Institute of Bioproduct Development (IBD), Faculty of Chemical Engineering, Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia

\*Corresponding author: i.liza@cheme.utm.my

## Article history


Received :1 October 2013

Received in revised form :

31 December 2013

Accepted :27 January 2014

## Graphical abstract



## Abstract

Currently, finding alternative ways of extracting medicinal plant gain more interest from the researchers. *Quercus infectoria*, a medicinal plant, is rich with bioactive compound being extracted using supercritical carbon dioxide (SC-CO<sub>2</sub>) extraction which helps to maintain the quality of the product as well as green environment. CO<sub>2</sub> is widely used as solvent due to moderate critical conditions, nontoxic and easily removed from the products. This work was performed to determine the optimum extraction parameters of SC-CO<sub>2</sub> extraction and their effects on the total phenolic content and antioxidant activity of *Q.infectoria* extract. Hence, two different parameters have been investigated which were extraction time and CO<sub>2</sub> flow rate (2, 3, 4 ml/min) while pressure (P) and temperature (T) were fixed at highest density (P = 30 MPa, T = 40°C). The results obtained from this study show that the solvent flow rate of 2 ml/min gives the highest percentage of yield (0.3652%) and the complete extraction of the sample was achieved at 80 minutes. Better quality of the extract was shown at 2 ml/min as resulted in high amount of phenolic compound in the extract presented as gallic acid equivalent (GAE) (2.04×10<sup>2</sup> mg GAE/g sample). The extracts were screened for possible antioxidant activity by 2,2-diphenyl-1-picryl hydrazyl (DPPH) free radical scavenging assays. In this study, the best result obtained for antioxidant activity was at flow rate of 3 ml/min with inhibition percentage of 96.97%.

**Keywords:** *Quercus infectoria*; SC-CO<sub>2</sub> extraction; total phenolic content; antioxidant activity

© 2014 Penerbit UTM Press. All rights reserved.

## 1.0 INTRODUCTION

Healing powers in medicinal plants have been used for centuries as remedies for human diseases and offer a new source of biologically active chemical compounds as antioxidant, antimicrobial agent and more, and it is clinically proven. Generally, over three-quarters of the world population depends generally on plants and plants extract for health care purposes [1]. This is due to the fact that medicinal plants have the richest bio-resources of drugs in traditional medicinal systems, modern medicines, nutraceuticals, folk medicines, food supplements, intermediate and chemical entitled for synthetic drugs [2]. There are three basic parameters influencing the quality of extract which are the plant part used as starting material, extraction solvent and thirdly is extraction procedure [3]. The researchers would keep on using the medicinal herb and these should be incorporated into modern medicine.

*Quercus infectoria*, known as manjakani is a type of medicinal plant that can be used in treating diseases and it is already well-known since ancient time. Oak (*Quercus*) is part of Fagaceae family. *Q.infectoria* galls is stimulated by gall-wasp called *Cynips quercufolii*, through laying its eggs in the bark. As a result, the inner cells of the bud becomes bigger [4]. During this time all starch will be converted into tannins, resulting in the galls containing tannins 60% from its total weight [5-6]. Besides, the galls also reported to restrain small amount of free gallic acid (2-4%), ellagic acid, syringic acid, β-sitosterol, amentoflavone, hexamethyl ether, isocryptomerin, methyl oleanate, methyl betulate and hexagalloyl glucose [7-10]. *Q.infectoria* was claimed to be extremely valuable for the local Malay postpartum women and there are no risky effect reported until now.

Until today, the application of supercritical fluid extraction (SFE) technology in industry has been limited, however the use of this technique already grab much attention for removal of organic

compounds from different liquid and solid matrices. Examples of SFE technology that being used are extraction of oil from fig leaf gourd seeds, extraction of bioactive compounds from *A.grossedentata* stems, extraction of essential oil from *Achillea santolina* and lycopene extraction from tomato peel [11-14]. SFE generally based on the utilization of a fluid under supercritical conditions, is a technology suitable for extraction and purification of a variety of compounds, especially for those that have low volatility or susceptible to thermal degradation. Supercritical fluids have higher diffusivity and lower density, surface tension and viscosity which can be varied by changing the operating conditions, subsequently give advantages to the extraction process. Furthermore, the use of CO<sub>2</sub> as a solvent in SFE can avoid the alteration of the product since the operating temperature used is near ambient temperature [15].

To the best of our knowledge, there is no research on the extraction of *Quercus infectoria* by supercritical fluid extraction using carbon dioxide (CO<sub>2</sub>) as solvent. Hence, a study has been conducted to determine the optimum extraction parameter (CO<sub>2</sub> flow rate, extraction time), total phenolic content and its antioxidant activity.

## 2.0 MATERIALS AND METHODS

### 2.1 Preparation of Plant Material

The galls of *Q.infectoria*, purchased from local market (Kota Tinggi, Johor Malaysia), were first separated, rinsed using tap water to remove unwanted material and dried in an oven at 60°C overnight. They were crushed by blender before extraction was done. The prepared seeds were stored in dark place at room temperature.

### 2.2 Supercritical Carbon Dioxide Extraction

The extraction of *Q.infectoria* galls was done using Supercritical Fluid Extraction Unit [16-17]. The system consists 50 ml extraction vessel, a high-pressure pump, an automated back pressure regulator and an oven. Liquid CO<sub>2</sub> was supplied from a gas cylinder.

The extraction was conducted using CO<sub>2</sub> flow rate of 2, 3 and 4 ml/min and the samples were taken for every 10 minutes. The pressure and temperature were fixed at 30 MPa and 40°C. At these conditions, the solvent is at highest density which is 0.919 g/ml.

### 2.3 DPPH Radical Scavenging Activity Assay

The assay of antioxidant activity was done using 2,2-diphenyl-1-picryl hydrazyl (DPPH) methanolic solution [18]. In this method, 77 µl of 2.5 mg/ml extracts and positive control (BHT) in methanol were added into 3 ml of methanolic DPPH solution. The mixture was vortex at room temperature for 30 seconds. The control sample absorbance ( $A_{\text{control}}$ ) which contains methanolic solution of DPPH was also measured. All of the mixtures were allowed to stand in a dark place for 30 minutes. The absorbance of all sample solutions was measured at 517 nm using UV-Vis spectrophotometer. The percentage of scavenging effect was calculated using Equation (1).

$$\% \text{ scavenging} = \frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \times 100 \quad (1)$$

### 2.4 TPC Determination


Total phenolic content (TPC) in the extracts was determined using Folin-Ciocalteu (FC) reagent [19]. Twenty µl of 1 mg/ml plant extract, 1.58 ml of distilled water and 100 µl of FC reagent (diluted

ten-fold) were mixed in a test tube. The mixture was left at room temperature in order to allow the reaction take place. After 7 minutes, 300 µl of 75 g/l sodium carbonate (Na<sub>2</sub>CO<sub>3</sub>) solution was added to the sample solution. The tube was kept in a dark place for 30 minutes at room temperature. The absorbance of the colour changes was measured at 765 nm. The calculation of TPC was done on the basis of the gallic acid standard curve which constructs using the same procedure and at concentrations of 0, 50, 100, 150, 250 and 500 mg/ml. The results were expressed as gallic acid equivalents (mg GAE/g extract sample).

## 3.0 RESULTS AND DISCUSSION

### 3.1 Effects of CO<sub>2</sub> Flow Rate and Extraction Time on the Extraction Yield

Figure 1 shows the effect of CO<sub>2</sub> flow rate at respective extraction time at pressure of 30 MPa and temperature of 40°C. The accumulative yield of the extract at CO<sub>2</sub> flow rate of 2 ml/min showed the highest value (0.37%) whereas the lowest percentage yield was given by 4 ml/min (0.29%).



**Figure 1** The effects of CO<sub>2</sub> flow rate and extraction time on the accumulative yield (%) of the extracts at P = 30 MPa and T = 40°C

At extraction time of 2 hours, the solute-solvent saturation was achieved. When the saturation occurs, the solubility is increasing as the residence time increased. Besides, extracting seeds with high oil content is best when lower solvent flow rate is used in order to hinder compaction of the sample in the vessel that may obstruct complete extraction of the oil [20]. In addition, when the flow rate of solvent increases, it enhance the cumulative yield of extract, but further increase of flow rate decrease the yield [21].

In order to determine the optimum extraction time, the experiment was done until the constant yield was obtained. Figure 1 shows that after 80 minutes, the accumulative yield was performed to be constant. Thus, the extraction time of 80 minutes was chosen to be optimum parameter due to less CO<sub>2</sub> consumption was utilized. Besides, time has no significant effect on extraction yield after maximum conversion was obtained [22].

### 3.2 Total Phenolic Content

Total phenolic content (TPC) of extracts were measured to investigate its contribution in antioxidant activity and to support the choice of parameters in term of solvent flow rate and extraction time. The phenol content in the extracts at 2, 3 and 4 ml/min was found to be 203.54 ± 10.56, 186.14 ± 7.46 and 193.61 ± 9.88 mg

GAE/g sample, respectively as shown in Table 1. Amount of phenol content when using CO<sub>2</sub> flow rate of 2 ml/min suggested that higher solubility of the sample in the solvent. When the solubility is high, the amount of desired compound extracted is increased and it can prevent impurities to be extracted.


**Table 1** Total phenolic content shows at different CO<sub>2</sub> flow rate at P = 30 MPa and T = 40°C

CO <sub>2</sub> flow rate (ml/min)	Total phenolic content (mg GAE/g sample)±SD
2	203.54 ± 10.56
3	186.14 ± 7.46
4	193.61 ± 9.88

SD: Standard deviation

### 3.3 Antioxidant activity of extract

The antioxidant activity assay was carried out to investigate the ability of *Q. infectoria* to scavenge free radicals in vitro by increasing the scavenging activity. Based on Figure 2, it shows that all of the extract obtained at the specified CO<sub>2</sub> flow rate give higher activity (%) than butylated hydroxyanisole (positive control) but the difference was insignificant (P>0.05). These results may explained by the fact that these extract enriched with phenolic compounds and comparable to positive control, which always play an important role in the antioxidant activity of the plant [23].



**Figure 2** The effect of CO<sub>2</sub> flow rate on the DPPH radical scavenging activity compared to positive control (BHT) at P = 30 MPa and T = 40°C

Figure 2 clearly demonstrate that the highest antioxidant activity was attained at CO<sub>2</sub> flow rate of 3 ml/min (96.965%) but the difference between flow rate of 3 ml/min and 2 ml/min is too low (0.04%). Hence, 2 ml/min of CO<sub>2</sub> flow rate is more favourable to be used as optimum extraction parameter due to positive results obtained from the analysis as well as low cost since less CO<sub>2</sub> was used.

### 4.0 CONCLUSION

Based on the yields of the *Q. infectoria* extracts using SC-CO<sub>2</sub> extraction at 30 MPa and 40°C, the highest yield was at CO<sub>2</sub> flow rate of 2 ml/min (0.37%) and the extraction time of 80 minutes. The total phenolic compound at the CO<sub>2</sub> flow rate is 2.04 × 10<sup>2</sup> mg GAE/g sample. All of the extracts possess high antioxidant activity and could act as natural antioxidants for human body.

### Acknowledgement

The authors gratefully acknowledge Minister of Higher Education (MOHE) for financial supports and Centre of Lipid Engineering and Applied Research (CLEAR), Universiti Teknologi Malaysia for the use of laboratory instruments/ equipments and their kind supports and research grant GUP (Q.J130000.7125.02H01) during this work.

### References

- Joy, P. P., J. Thomas, S. Mathew, and B. P. Skaria. 1998. *Medicinal Plants*. Aromatic and Medicinal Plants Research Station, Kerala Agricultural University, India.
- Hammer, K. A., C. F. Carson, and T. V. Riley. 1999. Antimicrobial Activity of Essential Oils and Other Plant Extracts. *Journal of Applied Microbiology*. 86(6): 985.
- Ncube, N. S., A. J. Afolayan, and A. I. Okoh. 2008. Assessment Techniques of Antimicrobial Properties of Natural Compounds of Plant Origin: Current Methods and Future Trends. *African Journal of Biotechnology*. 7(12): 1797.
- Cook, W. M. H. 1869. *The Physio-Medical Dispensary: A Treatise on Therapeutics, Materia Medica and Pharmacy. Accordance with the Principles of Physiological Medication*. Boulder, Colorado: W.M.H. Cook.
- Muhsin, A. 2003. *Photo Herbs dictionary*. Beirut, Lebanon: AI-A'jami printing Co.
- Mahmoud, D. 1979. *Classification of Forest Trees*. College of Agriculture and forests, Mousil University.
- Ikram, M., and F. Nowshad. 1977. Constituents of *Quercus infectoria*. *Planta Medica*. 31: 286.
- Wiat, C., and A. Kumar. 2001. *Practical Handbook of Pharmacognosy Malaysia*. Malaysia: Pearson Education Malaysia Sdn. Bhd.
- Bruneton, J. 1995. *Pharmacognosy, Phytochemistry, Medicinal Plants*. Lavoisier, Paris: 102.
- Hwang, J. K., T. W. Kong, N. I. Baek, and Y. R. Pyun. 2000. Alpha-glycosidase Inhibitory Activity of Hexagalloylglucose from the Galls of *Quercus infectoria*. *Planta Medica*. 66: 273.
- Bernardo-Gil, M. G., I. M. G., Lopes, M. Casquilho, M. A. Ribeiro, M. M. Esquível, and J. Empis. 2007. Supercritical Carbon Dioxide Extraction of a Corn Oil. *Journal of Supercritical Fluids*. 40: 344.
- Wang, Y., L. Ying, D. Sun, S. Zhang, Y. Zhu, and P. Xu. 2011. Supercritical Carbon Dioxide Extraction of Bioactive Compounds from *Ampelopsis grossedentata* Stems: Process Optimization and Antioxidant Activity. *International Journal of Molecular Sciences*. 12: 6856.
- Farajpour, M., M. Ebrahimi, R. Amiri, S.A.S. Nori, and R. Golzari. 2011. Investigation of Variations of the Essential Oil Content and Morphological Values in Yarrow (*Achillea Santolina*) from Iran. *Journal of Medicinal Plants Research*. 5(17): 4393.
- Machmudah, S., Zakaria, S. Winardi, M. Sasaki, M. Goto, N. Kusumoto, and K. Hayakawa. 2012. Lycopene Extraction from Tomato Peel By-product Containing Tomato Seed Using Supercritical Carbon Dioxide. *Journal of Food Engineering*. 108: 290.
- Perrut, M. 2004. *Application of Supercritical Fluid Solvents in the Pharmaceutical Industry. Ion Exchange and Solvent Extraction*. New York: Marcel Dekker, Inc.
- Mandana, B., A.R. Russly, G. Ali, and S. T. Farah. 2011. Antioxidant Activity of Spearmint (*Mentha spicata* L.) Leaves Extracts by Supercritical Carbon Dioxide (SC-CO<sub>2</sub>) Extraction. *International Food Research Journal*. 18: 543.
- Machmudah, S., M. Kondo, M. Sasaki, M. Goto, J. Munemasa, and M. Yamagata. 2008. Pressure effect in Supercritical CO<sub>2</sub> Extraction of Plant Seeds. *Journal of Supercritical Fluids*. 44: 301.
- Patra, J.K., A.D. Mohapatra, S.K. Rath, N.K. Dhal, and H. Thatoi. 2009. Screening of Antioxidant and Antifilarial Activity of Leaf Extracts of *Excoecaria agallocha* L. *International Journal of Integrative Biology*. 7(1): 9.
- Singleton, V. L., and J. A. Rossi. 1965. Colorimetry of Total Phenolics with Phosphomolybdic-phosphotungstic Acid Reagents. *American Journal of Enology and Viticulture*. 16: 144.
- King, J. 1997. Chapter 17: *Critical Fluids for Oil Extraction- Technology and Solvents for Extracting Oilseeds and Nonpetroleum Oils*. In Wan, P.J., and Wakelyn, P.J. (Eds.). AOCS Press-Champaign, IL: 283–310.
- Kumoro, A.C., and M. Hasan. 2006. Effect of Solvent Flow Rate on the Supercritical Carbon Dioxide Extraction of *Andrographolide* from *Andrographis paniculata*. 11<sup>th</sup> Asia Pacific Confederation of Chemical

- Engineering Conference (11<sup>th</sup> APPChE 2006). Kuala Lumpur Convention Centre (KLCC), Malaysia.
- [22] Liza, M.S., R. Abdul Rahman, B. Mandana, S. Jinap, A. Rahmat, I.S.M. Zaidul, and A. Hamid. 2010. Supercritical Carbon Dioxide Extraction of Bioactive Flavonoid from *Strobilanthes crispus* (Pecah Kaca). *Food and Bioproducts Processing*. 88: 319.
- [23] Pourmorad, F., S.J. Hosseinimehr, and N. Shahabimajd. 2006. Antioxidant Activity, Phenol and Flavonoid Contents of Some Selected Iranian Medicinal Plants. *African Journal of Biochemistry*. 5(11): 1142.