

## Response of *Streptococcus zooepidemicus* to Oxidative Stress in Hyaluronic Acid Fermentation

Mashitah, M.D.<sup>1\*</sup>, Masitah, H.<sup>2</sup> and Ramachandran, K.B.<sup>3</sup>

<sup>1</sup>School of Chemical Engineering, Engineering Campus, Universiti Sains Malaysia, Seri Ampangan, 14300 Nibong Tebal, Penang, Malaysia

<sup>2</sup>Department of Chemical Engineering, Faculty of Engineering, University of Malaya, 50603 Kuala Lumpur, Malaysia

<sup>3</sup>Department of Biotechnology, Indian Institute of Technology Madras, Chennai 600 036, India

\*E-mail: chmashitah@eng.usm.my

### ABSTRACT

*Streptococcus zooepidemicus* (SZ) is an aerotolerant bacteria and its ability to survive under reactive oxidant raises the question of the existence of a defense system against oxidative stress. As a characteristic of lactic acid bacteria, *Streptococcus* lacks an ordinary anti-oxidative stress enzyme, catalases and an electron transport chain. Whether this bacterium resists oxidative stress prior to an exposure to a higher level of an oxidizing agent H<sub>2</sub>O<sub>2</sub> in hyaluronic acid fermentation is not known. This paper describes that *Streptococcus* cells, once treated with lower concentrations of H<sub>2</sub>O<sub>2</sub> (i.e. 0.25, 0.50 and 1.0 mM) at least, were prepared for a subsequent higher concentrations of H<sub>2</sub>O<sub>2</sub> such as 20.5 and 100 mM. At low concentrations (i.e. 0.25, 0.50 and 1.0 mM), H<sub>2</sub>O<sub>2</sub> was found to act as a stimulant for HA synthesis, but it became toxic if presented at a very high level (100 mM H<sub>2</sub>O<sub>2</sub>). The highest HA yield to glucose consumed ( $Y_{HA^{total}/glu}$ ) was 0.017 gg<sup>-1</sup> for the cells pre-treated with 0 mM of H<sub>2</sub>O<sub>2</sub>, and then exposed to 20.5 mM H<sub>2</sub>O<sub>2</sub>. Thus, this implied that this bacteria might possess a defense mechanism against oxidative stress and that this system was inducible.

**Keywords:** Hyaluronic acid, hydrogen peroxide, oxidative stress, protective response, *Streptococcus zooepidemicus*

### ABBREVIATIONS

DNA	deoxy-ribonucleic acid
HA	hyaluronic acid
H <sub>2</sub> O <sub>2</sub>	hydrogen peroxide
OD	optical density
ROS	reactive oxygen species
SBA	sheep blood agar
sHA	soluble hyaluronic acid
tHA	total hyaluronic acid

---

Received: 29 May 2007

Accepted: 22 August 2008

\*Corresponding Author

$Y_{HAtotal/glu}$	hyaluronic acid yield to glucose consumed
$Y_{H2O2/glu}$	hydrogen peroxide yield to glucose consumed
$Y_{x/glu}$	biomass yield to glucose consumed

## INTRODUCTION

The ability to persist and thrive under oxidative stress is necessary for the growth of bacteria, under aerobic or reactive oxidant challenges. In bacteria, the association of enzymatic and non-enzymatic systems, with reactive oxygen species (ROS), provides a mechanism for a cellular response against deleterious effects of the molecules. Of the various reactive molecules ( $O_2^-$ ,  $H_2O_2$ ,  $\bullet OH$ ), hydrogen peroxide ( $H_2O_2$ ) is particularly derived from non-radical oxygen. Though  $H_2O_2$  is chemically less reactive, it is still a threat to the structure and function of cells (Halliwell and Gutteridge, 1989). Likewise, it can readily diffuse across the cellular membranes and oxidatively damage a number of vital cellular components, including membrane lipids, enzymes and DNA (Miller and Britigan, 1997; Porter, 1984).

An adaptation to hyperoxidative environment provides organisms with a wider selection of ecological niches for survival. Like many other lactic acid bacteria, most *Streptococcus* species have manganese form of superoxide dismutase (Mn-SOD) for detoxification of ROS (Jakubovics, Smith and Jenkinson, 2002). However, this might not be the only mechanism in bacteria. The binding of the metal ions onto superoxide dismutase (SOD) or catalases in forms, which was unable to accelerate radical ( $O_2^-$ ,  $H_2O_2$ ,  $\bullet OH$ ) or non-radical reactions (Halliwell and Gutteridge, 1999), was thought to form the basis for an alternative chemically detoxification of  $H_2O_2$ . These involved reactions are described as (Bannister, Bannister and Rotilio, 1987):  $2O_2^- + 2H^+ \rightarrow H_2O_2 + O_2$  (Superoxide dismutase),  $2(H_2O_2) \rightarrow 2H_2O + O_2$  (Catalase) and  $H_2O_2 + RH_2 \rightarrow 2H_2O + R$  (Non-specific peroxidase). Such a system might be necessary in reducing the levels of  $H_2O_2$  and damaging their production. However, as is the characteristic of lactic acid bacteria, *Streptococcus* species does not synthesize heme and lacks catalase (Condon, 1987). The specific mechanisms by which they adapt to peroxide stress are still unknown.

Traditionally, hyaluronic acid (HA), which is an industrially important biopolymer, was extracted from rooster comb. However, an increasing trend to streptococcal fermentations is rapidly expanding in view of its emerging applications in the medical and cosmetic industries (Lerner, 1996). Remarkly, *Streptococcus zooepidemicus* (SZ) were also greatly used as the risk of cross species viral infection which can be avoided (Huang *et al.*, 2006). Nevertheless, there were some drawbacks in relation to *Streptococcus* HA production routes. According to Goh (1998), under aerated conditions, HA production was found to be reduced as a result of the growth inhibition by hydrogen peroxide ( $H_2O_2$ ), since this compound is inherently toxic and reactive, as well as *Streptococcus zooepidemicus* being catalase negative (Hardie and Whiley, 1995). Mashitah *et al.* (2005) reported that  $H_2O_2$  produced by *Streptococcus zooepidemicus* cells, did not affect the growth of cell, but influenced the production of HA. Accordingly, the production of  $H_2O_2$  took place during the growth phase, and this was only started after the growth had reached its late exponential phase, that is, when  $H_2O_2$  in the culture media had depleted. Whether such cells are able to resist oxidative stress prior to exposure to higher level of  $H_2O_2$ , this has not been clearly defined. In this study, the researchers examined the response of *Streptococcus zooepidemicus* cells to oxidative stress, prior and during the HA fermentation. For this purpose,  $H_2O_2$  was used as an oxidizing agent.

## MATERIALS AND METHODS

### *Strain*

*Streptococcus equi* sub-species *zooepidemicus* ATCC 39920 was obtained from the American Type Culture Collection (Rockville, MD., USA). It was maintained on sheep blood agar (SBA) slants and kept at 4°C.

### *Culture medium*

The composition of the medium used in all the experiments comprised of (g<sup>l</sup><sup>-1</sup>) glucose 30, yeast extract 10, KH<sub>2</sub>PO<sub>4</sub> 0.5, Na<sub>2</sub>HPO<sub>4</sub>·12H<sub>2</sub>O 1.5, and MgSO<sub>4</sub>·7H<sub>2</sub>O 0.5, respectively. The pH of the medium was adjusted to 7.0 with 5 M NaOH prior to autoclaving it at 121°C for 20 min. This glucose solution was autoclaved separately and mixed aseptically with the other components on cooling.

### *Cell suspension*

Cell suspension of the shake flask culture was prepared by inoculating aseptically a stock culture of *S. zooepidemicus* onto Sheep Blood Agar (SBA)-plates and incubated overnight at 37°C. The formed colonies were punched by a sterile cork borer to obtain ten round disks of 0.85 cm in diameter. The disks were then put in a sampling bottle containing 50 ml of sterile distilled water. The sampling bottle was vortexed for 3 min so that the cells could evenly be distributed in the liquid.

### *The Effect of H<sub>2</sub>O<sub>2</sub> Pre-treatment*

20 SBA-disks full of *S. zooepidemicus* colonies, were pre-incubated with 100 ml of H<sub>2</sub>O<sub>2</sub> solution (0, 0.25, 0.50 and 1.0 mM) in an orbital shaker at 37°C, 150 rpm for 30 min. These cells (15 ml) were then re-treated with H<sub>2</sub>O<sub>2</sub> (15 ml) at the indicated concentrations (0, 20.5 and 100 mM) into a flask containing 120 ml culture media. The cells were left to be in contact with H<sub>2</sub>O<sub>2</sub> in an orbital shaker at 37°C, 250 rpm for 24 h. After 24 h, the samples were analyzed for cell biomass, H<sub>2</sub>O<sub>2</sub> and HA production, and glucose consumption. The cell biomass was taken as indices of the cells growth during the fermentation period.

### *Analytical Methods*

The cell concentration was determined by measuring the optical density (*OD*) at 600 nm by Jenway Spectrophotometer and dry cell method. A correlation between the dry cell weight and *OD*<sub>600</sub> was established. The concentration of H<sub>2</sub>O<sub>2</sub> was analyzed using the spectrophotometric method as suggested by Emiliani and Riera (1968), with a slight modification. Hyaluronic acid (HA) and glucose concentrations were determined using the method described by Mashitah *et al.* (2002), and the hexokinase method (Sigma Diagnostic, Glucose HK, Procedure No 16-UV), respectively. The HA soluble (*HA*<sub>sol</sub>) represented the hyaluronic acid (HA) which had been solubilised or released from *Streptococcus zooepidemicus* capsule during the fermentation. The HA total (*HA*<sub>total</sub>) represented the combination of HA which was solubilised from the capsule during the fermentation and the HA which was released from the ruptured capsule into the solution, after being treated with sodium dodecyl sulphate and vortexed for 3 mins.

## RESULTS AND DISCUSSION

### *The Effect of H<sub>2</sub>O<sub>2</sub> Pre-treatment on Growth and Biomass Yield*

The ability of *S. zooepidemicus* to respond to H<sub>2</sub>O<sub>2</sub> pre-treatment, prior to HA fermentation, was investigated and the results are shown in Fig. 1. It was suggested that prior to treating the *Streptococcal* cells directly with a high level of H<sub>2</sub>O<sub>2</sub> in the production medium, the cells were pre-treated in the absence or in the presence of 0.25, 0.50 and 1.0 mM H<sub>2</sub>O<sub>2</sub> at 37°C, 150 rpm for 30 mins. This was done to allow the cells to acclimatize with the low levels of H<sub>2</sub>O<sub>2</sub> before being introduced to a medium with higher H<sub>2</sub>O<sub>2</sub> concentrations, since exposing to higher doses of H<sub>2</sub>O<sub>2</sub> at the start of the culture would significantly decrease the growth and HA production (Mashitah *et al.*, 2005; Mashitah, 2006).

As shown in Fig. 1(a), the *S. zooepidemicus* biomass was found to slightly increase when pre-treated with lower doses of H<sub>2</sub>O<sub>2</sub>. For a 0 mM added H<sub>2</sub>O<sub>2</sub> with 1.0 mM H<sub>2</sub>O<sub>2</sub> pre-treatment, it led to a slight increase in biomass as compared to the one without any pre-treatment. As for 20.5 mM added H<sub>2</sub>O<sub>2</sub>, with 0.5 mM H<sub>2</sub>O<sub>2</sub> pre-treatment, a slightly lower level of cell biomass was detected. A similar trend was also observed for 0.25 mM pre-treated cells. However, it was also observed that with 100 mM H<sub>2</sub>O<sub>2</sub> medium, not more than 1% of either 0, 0.25, 0.50 or 1.0 mM pre-treated population could survive as compared to the non-pretreated control. This showed that the treated cells when exposed to lower levels of H<sub>2</sub>O<sub>2</sub>, were better able to cope with subsequent toxic doses. This also meant that the growth was unaffected or even enhanced by pre-treating the cells to H<sub>2</sub>O<sub>2</sub>. The treatment with the highest H<sub>2</sub>O<sub>2</sub> concentration (100 mM) led to a depletion of the biomass values. Thus, indicating that the organism might have been killed or destroyed at this level of added H<sub>2</sub>O<sub>2</sub>.

### *The Effect of H<sub>2</sub>O<sub>2</sub> Pre-treatment on HA and Extracellular H<sub>2</sub>O<sub>2</sub> Production and Glucose Consumption*

As observed in Fig. 1 (a, b, c, d and e), the biomass and HA production (HA total and HA soluble) were closely related. For the cells treated with 0 mM H<sub>2</sub>O<sub>2</sub>, lesser amounts of extracellular H<sub>2</sub>O<sub>2</sub> and glucose consumption were detected (Fig. 1(d)). This could be due to the fact that during H<sub>2</sub>O<sub>2</sub> exposure, *S. zooepidemicus* might consume H<sub>2</sub>O<sub>2</sub> and glucose during growth. Therefore, when the maximum biomass was attained, H<sub>2</sub>O<sub>2</sub> became limited, and hence the inhibition was reduced to increase the production of HA. This showed that some cells might have evolved highly efficient and often redundant repair mechanisms to remove lesion from DNA, proteins and membrane lipids. For example, all damages produced by free radical attack on DNA molecules were repaired by a universal DNA repair process known as the base-incision repair (Demple and Harrison, 1994; Thibessard *et al.*, 2001; Zhang, 2002), indicating that the appearance of ROS, H<sub>2</sub>O<sub>2</sub> had led to the development of defense mechanisms which either kept the concentration of the oxygen derived radicals at acceptable levels or repaired oxidative damages.

The findings of the current study also showed that the main physiological benefit of adaptive response was clear to protect *Streptococcus* cells from higher doses of a toxic agent. Such a protective response also indicated that the cell, once exposed to the H<sub>2</sub>O<sub>2</sub>, expects, or at least is prepared for a subsequent lethal dose. Besides that, a protective mechanism could have occurred; the cocoid cells associated into strands by the HA capsule, where the reduced surface-to-volume ratio and the limited H<sub>2</sub>O<sub>2</sub> diffusivity in the capsule shielded the cells from H<sub>2</sub>O<sub>2</sub>. In other words, the streptococcal cells synthesized HA excessively for a reduced rate of H<sub>2</sub>O<sub>2</sub> uptake.

Response of *Streptococcus zooepidemicus* to Oxidative Stress in Hyaluronic Acid Fermentation




Fig. 1: The effect of hydrogen peroxide pretreatment on the growth and HA production, H<sub>2</sub>O<sub>2</sub> released and glucose consumption by *S. zooepidemicus* after 24 hr of fermentation period. (X-axis - concentration of treated H<sub>2</sub>O<sub>2</sub>; legend - concentration of the pre-treated H<sub>2</sub>O<sub>2</sub>)

When pre-treated cells were treated with higher doses of H<sub>2</sub>O<sub>2</sub> (100 mM), the growth and HA production was negligible (Fig. 1). As a result, a lesser amount of glucose was utilized (Fig. 1 (e)) and lower extracellular H<sub>2</sub>O<sub>2</sub> was detected in the media. This also means that a significant damage, due to H<sub>2</sub>O<sub>2</sub>, indiscriminately reacted with various macromolecules in the cells had occurred, thus leading to a variety of biochemical and physiological lesions, and resulting in metabolic impairment and cell death (Totter, 1980; Harman, 1981; Ames, 1983).

From Table 1, it can be observed that the highest HA yield to glucose consumed ( $Y_{HA_{total}/glu}$ ) was  $0.017 \text{ gg}^{-1}$  for the cells pre-treated with 0 mM, and then exposed to 20.5 mM  $\text{H}_2\text{O}_2$ . As for the cells pre-treated with 0.50 mM and exposed to 20.5 mM  $\text{H}_2\text{O}_2$  medium, the detected amount of extracellular  $\text{H}_2\text{O}_2$  in the media was found to be the highest. The highest by-product yield ( $Y_{H2O2}/glu$ ) to glucose consumed was  $1.44 \text{ mM g}^{-1}$ . According to Ryan and Kleinberg (1975), the concentration of  $\text{H}_2\text{O}_2$  increased continuously during the growth due to the catabolic activity, and the categorization of this organism as  $\text{H}_2\text{O}_2$  producer was therefore appropriate.

TABLE 1  
Yield parameters of *S. zooepidemicus* at various pre-treated  $\text{H}_2\text{O}_2$  concentrations, which were re-treated with 20.5 mM  $\text{H}_2\text{O}_2$  medium (Condition: T =  $37^\circ\text{C}$ , agitation 250 rpm)

Pretreated $\text{H}_2\text{O}_2$ conc. (mM)	Biomass, X ( $\text{g L}^{-1}$ )	Hyaluronic acid total, $HA_{total}$ ( $\text{g L}^{-1}$ )	Hydrogen peroxide, $\text{H}_2\text{O}_2$ (mM)	$Y_{x}/glu$ ( $\text{gg}^{-1}$ )	$Y_{HA}/glu$ ( $\text{gg}^{-1}$ )	$Y_{H2O2}/glu$ ( $\text{mM g}^{-1}$ )
0	0.672	0.388	29.7	0.0299	0.017	1.32
0.25	0.716	0.334	27.2	0.0288	0.013	1.09
0.50	0.798	0.364	37.3	0.0308	0.014	1.44
1.0	0.781	0.326	6.99	0.0362	0.015	0.32

## CONCLUSIONS

In the response of *S. zooepidemicus* cells to oxidative stress in hyaluronic acid fermentation, several important features were found. Streptococcal cells were aerotolerant bacteria; when treated to lower levels of  $\text{H}_2\text{O}_2$  (0, 0.25, 0.5 and 1.0 mM) it was better able to cope with the subsequent higher toxic levels of the oxidizing agent (20.5 mM).  $\text{H}_2\text{O}_2$ , at low level, acts as a stimulant for the cells to synthesize HA. At higher level (100mM), the cells might be destroyed or killed. Furthermore, for the Streptococcal cells to survive against the oxidative stress, a defense system which is inducible must exist.

## ACKNOWLEDGEMENTS

The authors are grateful to the University Sains Malaysia, the Ministry of Higher Education (FRGS Grant) and the University of Malaya (Vote F: 0066/2001A, 0148/2002 & 0158/2003) for the financial support granted for this research.

## REFERENCES

- AMES, B.N. (1983) Dietary carcinogens and ant carcinogens. Oxygen radicals and degenerative diseases. *Science*, 221, 1256-1264.
- BANNISTER, J.W., BANNISTER, W.H. and ROTILIO, G. (1987). Aspects of structure, functions and applications of superoxide dismutase. *CRC. Critical Rev. Biochem.*, 22, 111-180.
- CONDON, S. (1987) Response of lactic acid bacteria to oxygen, *FEMS Microbiol. Rev.*, 46, 269-280.
- DEMPLE, B. and HARRISON, L. (1994). Repair of oxidative damage to DNA: Enzymology and biology. *Annu. Rev. Biochem.*, 63, 915-948.

Response of *Streptococcus zooepidemicus* to Oxidative Stress in Hyaluronic Acid Fermentation

- EMILIANI, E. and RIERA, B. (1968). Enzymatic oxalate decarboxylation in *Aspergillusniger*. II. Hydrogen peroxide formation and other characteristics of the oxalate decarboxylase. *Biochimica et Biophysica Acta*, 167, 414-421.
- GOH, L.T. (1998). Fermentation studies of hyaluronic acid production by *Streptococcus zooepidemicus*. (PhD Thesis. University of Queensland, Brisbane, Australia).
- HALLIWELL, B. and GUTTERIDGE, J.M.C. (1989). *Free Radicals in Biology and Medicine* (2<sup>nd</sup> Edition). Oxford: Clarendon Press.
- HALLIWELL, B., GUTTERIDGE, J.M.C. (1999). *Free Radicals in Biology and Medicine* (3<sup>rd</sup> Edition). Oxford: Oxford University Press.
- HARDIE, J.M. and WHILEY, R.A. (1995). The genus *Streptococcus*. In B.J.B. Wood and W.H. Holzapfel (Eds.), *The genera of lactic acid bacteria 2* (pp. 55-124). Blackie Academic and Professional.
- HARMAN, P. (1981). The aging process. *Proc. Natl. Acad. Sci.*, 78, 7124-7128.
- HUANG, W.C., CHEN, S.J. and CHEN, T.L. (2006). The role of dissolved oxygen and function of agitation in hyaluronic acid fermentation. *Biochem. Eng. J.*, 32, 239-243.
- JAKUBOVICS, N.S. and JENKINSON, H.F. (2001). Out of the iron age: new insights into the critical role of manganese homeostasis in bacteria. *Microbiology*, 14, 1709-1718.
- JAKUBOVICS, N.S., SMITH, A.W. and JENKINSON, H.F. (2002). Oxidative stress tolerance is manganese (Mn<sup>2+</sup>) regulated in *Streptococcus gordonii*. *Microbiology*, 148, 3255-3263.
- LENER, M. (1996). Hyaluronic acid market benefits from new uses. *Chemical Marketing Reporter*, 250, 12-13.
- MASHITAH, M.D., MASITAH, H. and RAMACHANDRAN, K.B. (2002). A rapid method for an identification of hyaluronic acid produced by fermentation of *Streptococcus zooepidemicus* ATCC 39920. In 15<sup>th</sup> Malaysian Analytical Chemistry Symposium and Regional Science Instrumentation Expo 2002, 10-12 September. Bayview Beach Resort, Pulau Pinang.
- MASHITAH, M.D., RAMACHANDRAN, K.B. and MASITAH, H. (2005). Sensitivity to hydrogen peroxide of growth and hyaluronic acid production by *Streptococcus zooepidemicus* ATCC 39920. *Dev. Chem. Eng. Mineral Process.*, 13(5/6), 1-12.
- MASHITAH, M.D. (2006). Kinetics of hydrogen peroxide formation and its relation to hyaluronic acid production by *Streptococcus zooepidemicus*. (PhD Thesis, University of Malaya, Kuala Lumpur, Malaysia).
- MILLER, R.A. and BRITIGAN, B.E. (1997). Role of oxidants in microbial physiology. *Clin. Microbiol. Rev.* 10, 1-18.
- PORTER, N.A. (1984). Chemistry of lipid peroxidation. *Methods Enzymol*, 105, 273-282.
- RYAN, C.S. and KLEINBERG, I. (1975) Bacteria in human mouths involved in the production and utilization of hydrogen peroxide. *Arch. Oral. Biol.*, 40(8), 753-763.
- THIBESSAND, A., FERNANDEZ, A., GITZ, B., BOURGET, N.L. and DECARIS, B. (2001). Hydrogen peroxide effects on *Streptococcus thermophilus* CNRZ 368 cell viability. *Res. Microbiol.* 152, 593-596.
- TOTTER, T.R. (1980). Spontaneous cancer and its possible relationship to oxygen metabolism. *Proc. Natl. Acad. Sci.* 79, 1763-1767.
- ZHANG, H. (2002). Function of oxygen-resistance proteins in the anaerobic bacteria *Desulforibrio vulgaris* strain Hildenborough. *MSc Thesis*, University of 1Calgary, Canada.